

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

# Spectra Simazine 900 WDG

## HERBICIDE

ACTIVE CONSTITUENT: 900g/kg SIMAZINE

GROUP **G** HERBICIDE

For the control of weeds in chickpeas, lupins, TT Canola, orchards, vineyards and certain other horticultural and non-crop situations as per the Directions for Use table.

**IMPORTANT: THIS LEAFLET IS PART OF THE LABEL ATTACHED TO THE CONTAINER. READ THOROUGHLY BEFORE OPENING OR USING THIS PRODUCT.**

Spectra Pacific (Australia) Pty Ltd ACN 127 711 479  
1/14 Palmer Place, Murarrie  
QLD 4172 AUSTRALIA  
Phone: 07 3907 0744 • Fax: 07 3890 4663  
APVMA Approval No.: 63399/0109

### DIRECTIONS FOR USE:

CROP / SITUATION	WEEDS CONTROLLED	STATE	RATE/ha		CRITICAL COMMENTS
			LIGHT SOIL	HEAVY SOIL	
Almonds	Amaranthus, Annual Ryegrass, Annual Thistles, Barley Grass, Bindy-eye, Capeweed,	SA only	1kg or 60g/100L water	2kg or 120g/100L water	Use only if trees are at least 3 years old. Apply to bare moist soil immediately after cultivation and before weeds emerge.
Asparagus	Chickweed, Common Sowthistle, Corn Gromwell (Sheepweed), Creeping Oxalis, Doublegee, Fat Hen, Fumitory, Ivy-Leaf Speedwell, Mustard, Native Geranium, Nettles, Potato Weed,	All States	1.3kg	2.5kg	Apply to bare moist soil after last cultivation and before spear emergence DO NOT apply to foliage or when fruit is present. Use on established plants only.
Berry Fruits	Fat Hen, Fumitory, Ivy-Leaf Speedwell, Mustard, Native Geranium, Nettles, Potato Weed,				Use only if trees are at least one year old.
Citrus	Red Shank, Shepherd's Purse, Turnip, Winter Grass, Wireweed (not Qld, Tas), Suppression of: Brome Grass, Soursob, Wild Oats	All States	1.2kg or 60g/100L water	1.2kg or 60g/100L water	Apply to bare moist soil after planting. High rates may cause crop damage on sandy soils low in organic matter.
Gladioli		Qld only	2kg	4kg	Use only if vines are at least two years old.
Grapevines		NSW, Vic, Tas, SA, WA only	1.25kg OR 60g/100L water	2.5kg OR 120g/100L water	Use only if vines are at least 12 months old. In first year of use split applications are preferred.
Hops		All States	1.3kg	2.5kg	Apply to moist weed free soil in late Winter, early Spring before hop emergence.
Canola (Triazine tolerant varieties only) Pre-emergence or post sowing pre-emergence only	Capeweed, Charlock, Clover, Corn Cromwell, Doublegee, Fumitories, Geraniums, Ivy-Leaf Speedwell, London Rocket, Mustards, Turnips, Paterson's Curse, Shepherd's Purse, Silver Grass (Vulpia). Suppression of Annual Ryegrass, Barley Grass, Brome Grass, Wild Oats and Wild Radish	All States	1.1kg	2.2kg	<b>Important: This use is subject to an INTEGRATED WEED MANAGEMENT STRATEGY TT Canola. See General Instructions: Integrate Weed Management Strategy for TT-Canola.</b> Can be applied up to a week before sowing or post-sowing pre-emergence (ideally incorporated by harrows). For best results, apply to bare moist soil, either immediately before seeding or as a pre-emergence treatment at or within 7 days of planting. Sufficient rainfall (20-30mm) to wet the soil through the weed root zone is necessary within 2-3 weeks of application. Application should not be made to ridged or excessively cloddy soil. When applied before seeding, incorporate to a depth of 5cm.
Chickpeas	Deadnettle, Indian Hedge Mustard, Lesser Swinecress, Milk Thistle, Prickly Lettuce, Purple Goosefoot, Shepherd's Purse, Turnip Weed, Wireweed and Suppression of Black Bindweed and Paradoxa Grass.	Qld, NSW, SA only	800g + 1.5L prometryne (500 g/L)		Apply immediately post-planting. Application should not be made to ridged or excessively cloddy soil. For reliable results significant rain, 20-30mm is necessary within 2-3 weeks of sowing.
	Capeweed, Clover, Dock, Doublegee, Mustard, Radish, Self Sown Cereals, Silver Grass, Turnip Weed and suppression of Barley Grass, Ryegrass and Wild Oats	WA only	550g	1.1kg	For best results, apply to bare moist soil, either immediately before seeding, or as a pre-emergence treatment at or within 7 days of planting. Sufficient rainfall (20-30mm) to wet the soil through the weed root zone is necessary within 2-3 weeks of application. Results can be variable if seasonal conditions are dry prior to sowing and chickpeas are sown into a dry or low moisture seedbed. Application should not be made to ridged or excessively cloddy soil. When applied before seeding, incorporation by the sowing operation should not be greater than 5cm. Use 0.55-0.775kg on lighter soils and in the Northern Agricultural areas and up to 1.1kg on heavier soil types. DO NOT use on whitish or grey sands.
Faba Beans	Annual Ryegrass, Barley Grass, Brome Grass, Capeweed, Corn Gromwell (Sheepweed), Deadnettle, Fumitories, Geranium, Ivy-leaf Speedwell, Medics, Mustards,	NSW, ACT, Vic, SA, WA only	1.1kg	1.4kg	Apply either pre-seeding or immediately post-sowing which is preferred on light soils. Sow the crop at least 5cm deep. Application should not be made to ridged or excessively cloddy soil. For fully reliable results significant rainfall, 20- 30mm is necessary within 2 to 3 weeks of sowing. DO NOT use rates higher than 2 litres per hectare on soils with pH 8.0 and above as unacceptable crop damage may occur.
	Paradoxa Grass, Saffron Thistle, Soursob, Volunteer Canola, Wireweed and suppression of Wild Oats		560g	830g	Apply to bare moist soil and incorporate to a depth of 5cm just prior to sowing. Incorporation should be made within 4 hours of application. Use this mixture where Annual Ryegrass and Wild Oats are the major problems. Application should not be made to ridged or excessively cloddy soil. For fully reliable results significant rainfall, 20-30mm is necessary within 2 to 3 weeks of application.
			Plus 800mL of trifluralin (480g/L)		

CROP / SITUATION (cont)	WEEDS CONTROLLED	STATE	RATE/ha		CRITICAL COMMENTS
			LIGHT SOIL	HEAVY SOIL	
Lupins	Amaranthus, Annual Ryegrass, Annual Thistles, Barley Grass, Bindy-eye, Capeweed, Chickweed, Common Sowthistle, Corn Gromwell (Sheepweed), Creeping Oxalis, Dou7blegee, Fat Hen, Fumitory, Ivy-Leaf Speedwell, Mustard, Native Geranium, Nettles, Potato Weed, Red Shank, Shepherd's Purse, Turnip, Winter Grass, Wireweed (not Qld, Tas)  Suppression of Brome Grass, Soursob, Wild Oats	Qld, NSW, Vic, Tas, SA only	1.3kg	2.2kg	Apply within 2 days of sowing. Seedbed should be moist and free of weeds, large clods and ridges. Apply higher rate for suppression of Wild Oats and Soursob. Use the lower rates (NSW, SA, Vic only) in combination with 8900mL of trifluralin (480g/L).
		NSW, Vic, SA only	830g	1.3kg	Apply anytime from 4 weeks before sowing until just before sowing. Apply as a tank mixture to a moist bare level soil and incorporate to a depth of 50mm prior to sowing. Incorporation must be within 4 hours of spraying especially under warm conditions. More effective than SIMAZINE on its own in dry conditions and where grass pressures are high. Use 830g/ha SIMAZINE on poorer light sands. 1.1kg/ha on sandy loams and on all other soil types. Increase the rate to 1.3kg/ha where Brome Grass is the major problem.
			Plus 1.25L of trifluralin (480g/L)		
		WA only	550g	1.1kg	DO NOT exceed 500g/ha on Lupin Variety Ultra. Apply immediately before seeding or within 6 days after sowing. Seed bed should be moist and free of weeds, large clods and ridges. Use 550g to 800g/ha on yellow sands, 1.1kg/a on other soil types. <b>Note:</b> Not for use on white or grey sands.
			550g	800g	Apply to bare moist soil just before sowing. Incorporate to a depth of 1-5cm within 4 hours of spraying. Soil should be free of large clods or ridges. Choose SIMAZINE rate according to comments in panel above. <b>Note:</b> Not for use on white or grey sands
Lupins When no weeds are present at time of sowing		WA only	550g		Apply 1-6 days before direct drilling the crop. Use the higher rate of SIMAZINE on weeds up to two weeks old and the lower rate of SIMAZINE with a high rate of knockdown herbicide on weeds more than 2 weeks old. DO NOT exceed 500g/ha on Lupin Variety Ultra. Note: Not for use on white or grey sands.
			Plus recommended rate of other knockdown herbicide		
Pome Fruit, Apples, Pears		Qld only	3kg OR 90g/100L water	4kg OR 120 g/100L water	Use only if trees are at least 2 years old. Apply to bare moist soil.
		NSW, Vic, Tas, SA, WA only	1.8kg/ha OR 90g/100L water	2.5kg/ha OR 120g/100L water	Use only if trees are at least 2 years old. Apply to bare moist soil.
Roses	Amaranthus, Annual Ryegrass, Annual Thistles, Barley Grass, Bindy-eye, Capeweed,	NSW, Vic, Tas, SA, WA only	1.3kg OR 60g/100L water	2.5kg OR 120g/100L water	Apply to bare moist soil. DO NOT use on plants less than 1 year old.
Strawberries	Chickweed, Common Sowthistle, Corn Gromwell (Sheepweed), Creeping Oxalis, Doublegee, Fat Hen, Fumitory, Ivy-Leaf, Speedwell, Mustard, Native Geranium, Nettles, Potato Weed, Red shank, Shepherd's Purse, Turnip, Winter Grass, Wireweed (not Qld, Tas) Suppression of Brome Grass, Soursob, Wild Oats	All States	1.2kg	1.2kg	Apply between polythene covered beds. DO NOT use on white or grey sands in WA.

CROP / SITUATION	WEEDS CONTROLLED	STATE	RATE/ha	CRITICAL COMMENTS
Sub Clover established lucerne and perennial grass pastures	Vulpia (Silvergrass), Rat's Tail Fescue, Squirrel-tail Fescue, Sand Fescue	NSW, Vic, SA only	550 - 900g	Apply 6 to 10 weeks after emergence of the pasture, but not before the third trifoliolate leaf stage of Sub Clover. Best results are obtained from application to young, well grazed, and actively growing plants. Some damage to Sub Clover may occur especially at the higher rates. DO NOT apply to sandy soils and use the lower rates on light textured soils. Apply under moist soil conditions; rain following application enhances results. DO NOT apply to stressed plants. Insect infestations should be controlled and Sub Clover allowed to recover before Spectra Simazine 900 WDG Herbicide is applied. DO NOT use herbicides for broadleaved weed control within 3 weeks of application.
		NSW, Vic only	550 - 800g plus 100 – 160mL of paraquat (250 g/L)	Apply 6 to 10 weeks after emergence of the pasture, but not before the third Trifoliolate leaf stage of Sub Clover. Best results are obtained from application to young, well grazed and actively growing plants. However the addition of paraquat improves the control of well-established plants. Add a non-ionic surfactant at 0.2% v/v (200mL per 100L). Under conditions of good soil moisture, control of other grasses and some broadleaved weed seedlings may occur. DO NOT apply to sandy soils and use the lower rates on light textured soils. Apply under moist conditions; rain following application enhances results. DO NOT apply to stressed plants. Insect infestations should be controlled and Sub Clover allowed to recover before the SIMAZINE 900 WDG plus Paraquat (250g/L) is applied. DO NOT use herbicides for broadleaved weed control within 3 weeks of application.
		SA only	400 – 550g plus 100 – 160mL of Paraquat (250g/L)	
	Vulpia (Silvergrass), Rat's Tail Fescue	WA only	400 – 500g	Apply within 8 weeks of emergence of Sub Clover and grasses. Under conditions of good soil moisture control of other grasses and some broadleaf weeds may occur. Ensure that there is a good stand of Subterranean Clover present before applying. Use the lower rate on light textured soils. DO NOT use with broadleaf weed herbicides within 3 weeks of using SIMAZINE 900 WDG. DO NOT tank mix with other herbicides or add crop oils or wetting agents. DO NOT use on Medics, or Red or White Clover. DO NOT overlap when spraying, otherwise damage may be observed.
CROP / SITUATION	WEEDS CONTROLLED	STATE	RATE/ha	CRITICAL COMMENTS
<b>Forestry</b> including farm tree plantations, <i>Pinus</i> , <i>Araucaria</i> , <i>Corymbia</i> and <i>Eucalyptus</i> tree plantings pre and post-planting (see General Instructions for further information on tree species)	Pre-emergence control of many annual and some perennial grasses and broadleaved weeds including: Amaranthus, Annual (Wimmera), Ryegrass, barley Grass, Barnyard Grass, Bindi, Brome Grass, Capeweed, Chickweed, Common Sowthistle, Corn Gromwell, (Iron/Sheepweed), Creeping Oxalis, Fat Hen, Fumitories, Geranium, Ivy-Leaf, Speedwell, Mustards, Native Geranium, Nettles, Potato Weed, Powell's Amaranth, Red Root Amaranth, Red Shank, Shepherd's Purse, Summer Grass, Thistles – Annual, Soldier, Variegated, Turnips, Turnip Weed, Wild Mustard, Wild Radish, Wild Turnip, Yellow Wood Sorrel, Weed suppression at higher rates Soursob, Wild Oats  Weeds suppressed WA Annual Ryegrass, Brome Grass, Dock, Double Gee, Radish, Sorrel, Soursob, Wild Turnip, Wireweed	All States	1.6 – 6.7kg/ha	SIMAZINE 900 WDG may be applied prior to or after planting. For most situations use 5.0 – 5.5kg/ha. On gravelly coarse textured soils use 2.8 to 4.0kg/ha. In WA and SA in medium rainfall zones (400-600mm) use 3.0 to 4.0kg/ha and for sandy low organic soils use 1.6 to 3.0kg/ha. On sandy soils allow 2 weeks and at least 50mm of rain between spraying and planting. Use a maximum rate of 1.6kg/ha for over-spraying eucalyptus plantations on sandy, low organic matter soils in WA & SA. For short term weed control or when applied with other pre-emergent herbicides use half rates according to the situation. Apply to bare ground. Best results will be achieved if applied to moist, finely tilled soil. If weeds are present add a knockdown or other partner herbicide as a tank mix. If spraying after planting check tree tolerance to knockdown or other partner herbicide before using. Application in a band 1.5 to 2.0m wide, over the planting row is recommended although broadcast application can also be used. Rates above refer to sprayed hectares and not field hectares. Some tree damage may occur especially with post-plant application. Recovery is normally rapid. Directed application is preferred for post-planting application especially when the higher rates are used.

Situation & Crops	Weeds Controlled	State	Rate kg/ha		CRITICAL COMMENTS
			Light Soil	Heavy Soil	
Non-crops uses, Commercial, Industrial, Rights of Way, Public Utility areas, Road shoulders, Drains, Headlands, Driveways, Railway tracks, Aerodromes, Gutters, Footpaths	As Above and including Barnyard Grass, Summer Grass	All States	5kg OR 500g/100L water	5kg OR 500g/100L water	Provides seasonal control of annual weed germinations. Apply to bare moist ground.
			10kg OR 1kg/100L water	24kg OR 2.4kg/100L water	Provides long residual control in Winter rainfall areas. Apply to bare moist ground.
			24kg OR 2.4kg/100L water	48kg OR 4.8kg/100L water	Provides long residual control in Summer rainfall areas. Apply to bare moist ground. Read remarks under Crop Safety.
Dams, Tanks, Troughs	Filamentous Blue-Green Algae	WA only	2g/1000 litres of water		Mix in a convenient amount of water and apply when algae development is first noticed.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

**WITHHOLDING PERIOD:**

***HARVEST:***

**ALL CROPS: NOT REQUIRED WHEN USED AS DIRECTED**

***GRAZING:***

**CANOLA: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 15 WEEKS AFTER APPLICATION**

**SUB CLOVER: DO NOT GRAZE FOR 14 DAYS AFTER APPLICATION**  
**DO NOT CUT FOR STOCK FOOD FOR 21 DAYS AFTER APPLICATION**

**CHICKPEAS: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 9 WEEKS AFTER APPLICATION**

**FABA BEANS: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 8 WEEKS AFTER APPLICATION**

**GENERAL INSTRUCTIONS**

SPECTRA SIMAZINE 900 WDG Herbicide is a pre-emergent herbicide, which is most effective when applied to bare, moist soil. Because it acts mainly through the roots of weeds, best results are obtained when rain or irrigation water move the herbicide down to the root zone within 2 weeks of application. The duration and effectiveness of control depends on the amount applied, the type of soil, intensity of leaching and the species of weeds present.

**RESISTANT WEEDS WARNING**

**GROUP C HERBICIDE**

SPECTRA SIMAZINE 900 WDG Herbicide is a member of the Triazine group of herbicides. SIMAZINE 900 WDG has the inhibitor photosynthesis mode of action. For weed resistance management SIMAZINE 900 WDG is a Group C herbicide. Some naturally occurring weed biotypes resistant to SIMAZINE 900 WDG Herbicide and other Group C herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by SIMAZINE 900 WDG Herbicide or Group C herbicides.

However, resistance should not be assumed without first viewing the method of application, the coverage and the timing of application. Resistance to SIMAZINE 900WDG cannot be predicted and may occur at any time. It is not due to a fault in the product and Spectra Pacific (Australia) Pty Ltd accepts no liability for any losses that may result from the failure of SIMAZINE 900 WDG Herbicide to control resistant weeds.

**INTEGRATED WEED MANAGEMENT STRATEGY FOR TT CANOLA**

An Integrated Weed Management Strategy for TT canola (the Strategy) has been developed and funded by the Canola Association of Australia in cooperation with certain companies.

The Strategy outlines recommendations, measures and options for weed management, including management of herbicide resistance in weed populations.

The Strategy is available from Spectra Pacific (Australia) Pty Ltd and the Canola

Rate kg/ha	CRITICAL COMMENTS	
	Light Soil	Heavy Soil
5kg OR 500g/100L water	5kg OR 500g/100L water	Provides seasonal control of annual weed germinations. Apply to bare moist ground.
10kg OR 1kg/100L water	24kg OR 2.4kg/100L water	Provides long residual control in Winter rainfall areas. Apply to bare moist ground.
24kg OR 2.4kg/100L water	48kg OR 4.8kg/100L water	Provides long residual control in Summer rainfall areas. Apply to bare moist ground. Read remarks under Crop Safety.
2g/1000 litres of water		Mix in a convenient amount of water and apply when algae development is first noticed.

Association of Australia. A program has been developed that outlines sound agronomic practices and integrated weed management programs designed to optimise the performance of TT Canola.

It is advised that consultation of IWM be undertaken with an accredited agronomist prior to use of SIMAZINE 900 WDG on TT Canola.

**TO MINIMISE HERBICIDE RESISTANCE**

- Avoid dry sowing in heavily weed-infested paddocks. Wait for weed germination after the opening rains in weedy paddocks. Use a pre-plant knockdown or cultivation. No weeds should be allowed to survive at this stage.
- Adapt the weed control program to the anticipated weed spectrum and pressure:

**Broadleaf Weeds and Ryegrass:** Use SIMAZINE 900 WDG or ATRAZINE 900 WDG plus trifluralin pre-emergence. A follow-up with a Group A herbicide (if ryegrass is susceptible) or ATRAZINE 900 WDG may be necessary.

**Broadleaf Weeds only:** Use ATRAZINE 900 WDG post-emergence.
- DO NOT use ATRAZINE 900 WDG or SIMAZINE 900 WDG if the area to be treated had a triazine herbicide applied to it last season.
- Watch for escapes, especially in paddocks with a long history of Group C herbicide use.
- DO NOT use Group C herbicides in consecutive years.

**TO AVOID TRIAZINE CARRY OVER**

**On acid soils (pH less than 6.5)** – The maximum rate of ATRAZINE 900 WDG or SIMAZINE 900 WDG or a combination of the two products to be applied to the crop during the growing season is 2.2kg/ha.

**On alkaline soils (pH greater than 6.5)** – The maximum rate of ATRAZINE 900 WDG or SIMAZINE 90 WDG or a combination of the two products to be applied to the crop during the growing season is 1.1kg/ha.

**Post-emergence use** – it is recommended that ATRAZINE 900 WDG only be used and at rates of 1.1 kg/ha or less, on both acid or alkaline soils.

**RESISTANT WEEDS REPORTING**

Growers should collect plant or seed samples where weeds that are normally susceptible to atrazine and simazine may be resistant, get them tested and seek professional advice.

**MIXING**

Half fill tank with clean water. Commence vigorous agitation. Add granules steadily. DO NOT dump. Ensure complete dispersion before filling to volume with water. Continue agitation during spraying. DO NOT leave spray solution in tank overnight.

**APPLICATION**

Use a boom sprayer or knapsack for small areas, 50 – 200L of mixture per hectare should be applied with the boom and a minimum equivalent to 500L per hectare through the knapsack sprayer. Clean spray equipment immediately after use using several flushings of water with detergent.

**TT CANOLA: APPLICATION**

DO NOT apply to TT Canola by aircraft. Apply only with a low boom sprayer with a 60m buffer zone downwind of treated fields to natural or impounded lakes or dams and a 20m buffer zone for any well, sink hole, intermittent or perennial stream. Apply only to areas where run-off is unlikely to occur or where run-off may be captured by farm earthworks.

**COMPATIBILITY**

This product may be mixed with amitrole, atrazine, diuron, glyphosate, diquat, paraquat and paraquat-diquat mixtures. Because the products may vary from time to time, a small scale compatibility test should be carried out before mixing in the spray tank. For residual control of annual weeds in non-crop areas, pome and citrus orchards and vineyards only, mixtures of the product and glyphosate may be used. Half fill the spray tank with clean water before adding this product at the recommended rate and mix thoroughly. Then add a 360g/L glyphosate product at 3L/ha or 700mL/100L and the remaining water. Maintain adequate agitation and use the mixture promptly. DO NOT use lower rates of glyphosate (360g/L) where control of perennial weeds is desired.

**PRECAUTION**

***Re-entry period:*** DO NOT enter treated areas without protective clothing until spray has dried.

**PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS**

DO NOT spray foliage of desired plants, except lupins (WA only). DO NOT use under peach, apricot, plus trees, ornamental, prunus or near newly planted shrubs.

DO NOT plant crops other than those recommended for at least nine months following treatments at rates up to 2.5kg/ha. At higher rates planting may not be possible for very long periods afterwards.

DO NOT use on sandy or porous soils near desirable plants or trees or where roots of such may extend or are near the surface. Crops of such as red beet, lettuce, cabbage, tomato, cucumber and carrots are particularly sensitive to simazine residues. DO NOT exceed highest rate recommended for any particular crop. DO NOT use on white or grey sands (WA only). DO NOT use in channels or drains where the roots of desirable plants may extend. DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

***Araucaria, Corymbis, Eucalyptus*** and ***Pinus*** plantations: Field trial and use experience has indicated that the following species are tolerant. Test tolerance before treating other species.

***Araucaria*** - all commercial species.
***Corymbia*** – *maculate*.
***Eucalyptus*** - *agglomerate, calophylla, camaldulensis, cladocalyx, cloeziana, dunnii, globules, grandis, laevopinea, nitens, pilularis, pseudoglobulus, regnans, rudis, saligna, wandoo*.
***Pinus*** - *caribaea* and hybrids, *ellotti, pinaster, radiata*

**SOIL MOISTURE**

If sufficient rain to thoroughly wet the weed root zone has not fallen within 14 days of application water in with the equivalent of 13mm of rain. If applied under dry conditions and irrigation is not available, mechanical incorporation to a depth of not more than 4cm within 7 days of sowing may assist results.

**PRODUCTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

DO NOT contaminate streams, rivers or streams with pesticide or used containers. DO NOT use the container for any other purpose. This product is very highly toxic to algae and aquatic macrophytes.

**STORAGE AND DISPOSAL**

**KEEP OUT OF REACH OF CHILDREN.** Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Shake empty bag into spray tank. DO NOT dispose of undiluted chemicals on site. Puncture or shred and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**SAFETY DIRECTIONS**

Avoid contact with eyes and skin. DO NOT inhale dust or spray mist. Wash hands after use.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia: 13 11 26)

**MATERIAL SAFETY DATA SHEET**

Additional information is listed in the Material Safety Data Sheet (MSDS). A material safety data sheet for SIMAZINE 900 WDG is available from Spectra Pacific (Australia) Pty Ltd upon request.

**CONDITIONS OF SALE**

The use of SPECTRA SIMAZINE 900 WDG Herbicide being beyond the control of the manufacturer, no warranty expressed or implied is given by Spectra Pacific (Australia) Pty Ltd regarding its suitability, fitness or efficacy for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Spectra Pacific (Australia) Pty Ltd accepts no responsibility for any consequence whatsoever resulting from the use of this product.